



Channels

Publication details, including instructions for authors and subscription information:
<http://www.tandfonline.com/loi/kchl20>

The unusual suspects: Regulation of retinal calcium channels by somatostatin

Michael E Hildebrand^a & Terrance P Snutch^b

^a Department of Neuroscience; Carleton University; Ottawa, ON Canada

^b Michael Smith Laboratories and Djavad Mowafaghian Centre for Brain Health; University of British Columbia; Vancouver, BC Canada

Accepted author version posted online: 25 Feb 2015.



[Click for updates](#)

To cite this article: Michael E Hildebrand & Terrance P Snutch (2015) The unusual suspects: Regulation of retinal calcium channels by somatostatin, Channels, 9:2, 61-62, DOI: [10.1080/19336950.2015.1018000](https://doi.org/10.1080/19336950.2015.1018000)

To link to this article: <http://dx.doi.org/10.1080/19336950.2015.1018000>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Versions of published Taylor & Francis and Routledge Open articles and Taylor & Francis and Routledge Open Select articles posted to institutional or subject repositories or any other third-party website are without warranty from Taylor & Francis of any kind, either expressed or implied, including, but not limited to, warranties of merchantability, fitness for a particular purpose, or non-infringement. Any opinions and views expressed in this article are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor & Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

It is essential that you check the license status of any given Open and Open Select article to confirm conditions of access and use.

The unusual suspects: Regulation of retinal calcium channels by somatostatin

Comment on: Farrell SR, et al. Somatostatin receptor subtype 4 modulates L-type calcium channels via $G\beta\gamma$ and PKC signaling in rat retinal ganglion cells. *Channels* 2014; 8(6):519-27; PMID:25483286; <http://dx.doi.org/10.4161/19336950.2014.967623>

Michael E Hildebrand^{1*} and Terrance P Snutch²; ¹Department of Neuroscience; Carleton University; Ottawa, ON Canada; ²Michael Smith Laboratories and Djavad Mowafaghian Centre for Brain Health; University of British Columbia; Vancouver, BC Canada; *Correspondence to: Michael E Hildebrand; Email: mike.hildebrand@carleton.ca; Keywords: calcium channel, excitotoxicity, $G\beta\gamma$, L-type, retinal ganglion cells, PKC, somatostatin receptor, inhibition; <http://dx.doi.org/10.1080/19336950.2015.1018000>

Excessive calcium influx is a major driver of retinal cell death and a hallmark of pathological processes triggered by retinal injury and disease.¹ Finding new ways to stop aberrant calcium entry is imperative in order to develop better therapeutics for preventing visual impairment and blindness. This effort requires an important balance between targeting the pathological functions of the players involved in calcium-mediated excitotoxicity while ensuring that their physiological functions remain intact.

The L-type class of voltage-gated calcium channels is one of the culprits implicated in calcium-mediated excitotoxicity. An increase in synaptic excitation and neuronal depolarization can lead to enhanced activation of L-type channels, prolonged inward calcium currents and an accumulation of intracellular calcium ions. In retinal ganglion cells (RGCs) – the output neurons of the retina – L-type channels have been shown to be the only calcium channel subtype to mediate calcium currents in both the soma as well as in RGC axons.² From this recent finding, it is tempting to speculate that blocking L-type channels in RGCs could be an effective therapeutic approach for preventing calcium-mediated degeneration in both the retina and optic nerve.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited. The moral rights of the named author(s) have been asserted.

However, generally blocking the L-type calcium channels themselves will likely interfere with numerous L-type-dependent physiological processes, ranging from pacemaker activity to gene expression to secretion.

An alternative therapeutic strategy for treating excitotoxic disorders is to identify individual modulators of L-type calcium channels which would be predicted to target pathological processes with increased specificity. The neuropeptide somatostatin is a key regulator of neuronal excitability in the retina. All five subtypes of G-protein coupled receptors for somatostatin (sst_1-5) are expressed in the retina, and there are well-established roles for sst_1 - and sst_2 -mediated signaling in visual processing and retinal homeostasis.¹ In contrast, the sst_4 receptor has been the neglected child of the somatostatin receptor family, with little known concerning its physiological relevance.³ In a recent impactful study, the laboratories of Steven Barnes and Nicholas Brecha demonstrated that the sst_4 subtype of somatostatin receptor is abundantly and specifically expressed in RGCs.⁴ This study also showed for the first time that activation of sst_4 selectively inhibits the L-type class of voltage-gated calcium channels in RGCs.

In this issue of *Channels*,⁵ Farrell and colleagues build upon their original study in neonatal RGCs to investigate modulation of L-type calcium channels by sst_4 in adult RGCs. Through this, they find that activation of sst_4 with the agonist L-803,087 inhibits calcium channel currents across RGC development. While a remaining caveat is that the selective inhibition of only L-type channels by sst_4

remains to be confirmed in acutely dissociated adult RGCs, the similarities found between the neonatal and adult cells strongly suggest developmental continuity. The unexpected twist in this study is the *unusual suspects* identified as mediating the inhibition of calcium channels via sst_4 .⁵ The authors show that in RGCs, calcium channel inhibition does not occur through the canonical $G\alpha_{i/o}$ -mediated signaling pathway typically downstream of sst_4 activation.³ Rather, they propose that sst_4 couples to the pertussis toxin-insensitive $G\alpha_q$ in RGCs. In support, they find that blocking PKC – which is downstream of $G\alpha_q$ activation – reduces the inhibition of calcium channels by L-808,087. Along with the contribution of PKC to the sst_4 -mediated inhibition of calcium currents in RGCs, the authors show that $G\beta\gamma$ is also required for a portion of the calcium current inhibition by L-803,087. Inhibition of voltage-gated calcium channels by $G\beta\gamma$ is typically voltage-dependent and reversed by strong depolarizing prepulses,⁶ yet Farrell and colleagues find that all sst_4 -mediated inhibition of calcium currents is voltage-independent. Finally, the effects of $G\beta\gamma$ - and PKC-mediated signaling on calcium channels are additive: blocking both signaling pathways together completely abolishes the sst_4 -mediated inhibition of calcium currents. Thus, in RGCs, the sst_4 receptor appears to activate distinct $G\alpha_q$ / PKC- and $G\beta\gamma$ - dependent signaling pathways, which converge to mediate the inhibition of putative L-type calcium channels.

Tapping into this unique endogenous sst_4 system may be an effective therapeutic strategy to selectively prevent calcium-mediated

excitotoxicity in RGCs, while leaving physiological sst_1 - and sst_2 -dependent processes unaffected. However, before this can occur the precise molecular events which couple sst_4 activation to L-type channel inhibition in RGCs need to be further understood. For example, which PKC isoform(s) are involved? Does the voltage-independent inhibition by $G\beta\gamma$ involve a direct physical interaction with calcium channels or, alternatively, is it through a secondary mediator such as the activation of potassium channels by $G\beta\gamma$? While there are no doubt many questions remaining, the findings of Farrell and colleagues⁵ represent a promising first step toward possible new

treatment avenues for retinal trauma and disease. Beyond the eye, both sst_4 and voltage-gated calcium channels are involved in gating pain signaling in peripheral sensory neurons.^{7,8} Thus, understanding mechanisms of coupling between sst_4 and calcium channels could also lead to new targets relevant to pain therapeutics.

References

1. Cervia D, et al. *Mol Cell Endocrinol* 2008; 286:112-22; PMID:18242820; <http://dx.doi.org/10.1016/j.mce.2007.12.009>
2. Sargoy A, et al. *PLOS One* 2014; 9:e84507; PMID:24416240; <http://dx.doi.org/10.1371/journal.pone.0084507>
3. Theodoropoulou M, Stalla GK. *Front Neuroendocrinol* 2013; 34:228-52; PMID:23872332; <http://dx.doi.org/10.1016/j.yfrne.2013.07.005>
4. Farrell SR, et al. *J Neurophysiol* 2010; 104:1347-54; PMID:20573967; <http://dx.doi.org/10.1152/jn.00098.2010>
5. Farrell SR, et al. *Channels* 2014; 8(6):519-27; PMID:25483286; <http://dx.doi.org/10.4161/19336950.2014.967623>
6. Zamponi GW, Snutch TP. *Proc Natl Acad Sci U S A* 1998; 95:4035-9; PMID:9520488; <http://dx.doi.org/10.1073/pnas.95.7.4035>
7. Gorham L, et al. *Eur J Pharmacol* 2014; 736:101-6; PMID:24769416; <http://dx.doi.org/10.1016/j.ejphar.2014.04.016>
8. Bourinet E, et al. *Physiol Rev* 2014; 94:81-140; PMID:24382884; <http://dx.doi.org/10.1152/physrev.00023.2013>